

Near-infrared survey of High Mass X-ray Binary candidates

J.M. Torrejón^{1,2}, I. Negueruela¹, D. M. Smith³, and T. E. Harrison⁴

¹ Departamento de Física, Ingeniería de Sistemas y Teoría de la Señal, Universidad de Alicante, E-03080 Alicante, Spain
e-mail: jmt@ua.es

² Massachusetts Institute of Technology, Kavli Institute for Astrophysics and Space Research, Cambridge MA 02139

³ Physics Department and Santa Cruz Institute for Particle Physics, University of California, Santa Cruz, 1156 High St., Santa Cruz, CA 95064

⁴ Astronomy Department, New Mexico State University, Box 30001/Department 4500, Las Cruces, NM 88003

Received ; accepted

ABSTRACT

Context. The *INTEGRAL* satellite is discovering a large population of new X-ray sources which were missed by previous missions due to high obscuration and, in some cases, very short duty cycles. The nature of these sources must be addressed via the characterization of their optical and/or infrared counterparts.

Aims. We investigate the nature of the optical counterparts to five of these newly discovered X-ray sources.

Methods. We combine infrared spectra in the *I*, *J*, *H* and *K* bands together with *JHK* photometry to characterize the spectral type, luminosity class and distance to the infrared counterparts to these systems. For IGR J19140+0951, we present spectroscopy from the red to the *K* band and new red and infrared photometry. For SAX J18186–1703 and IGR J18483–0311, we present the first intermediate-resolution spectroscopy reported. Finally, for IGR J18027–2016, we present new *I* and *K* band spectra.

Results. We find that four systems harbour early-type B supergiants. All of them are heavily obscured, with $E(B - V)$ ranging between 3 and 5, implying visual extinctions of ~ 9 to 15 magnitudes. We refine the published classifications of IGR J18027–2016 and IGR J19140+0951 by constraining their luminosity class. In the first case, we confirm the supergiant nature and rule out class III. In the second case, we propose a slightly higher luminosity class (Ia instead of Iab) and give an improved value of the distance based on new optical photometry. Two other systems, SAX J18186–1703 and IGR J18483–0311 are classified as Supergiant Fast X-ray Transients (SFXTs). XTE J1901+014, on the other hand, contains no bright infrared source in its error circle.

Conclusions. Owing to their infrared and X-ray characteristics, IGR J18027–2016 and IGR J19140+0951, emerge as Supergiant X-ray binaries with X-ray luminosities of the order of $L_X \sim [1 - 2] \times 10^{36} \text{ erg s}^{-1}$, while SAX J1818.6–1703 and IGR J18483–0311, turn out to be SFXTs at 2 and 3 kpc, respectively. Finally, XTE J1901+014 emerges as a puzzling source: its X-ray behaviour is strongly reminiscent of the SFXTs but a supergiant nature is firmly ruled out for the counterpart. We discuss several alternative scenarios to explain its behaviour.

Key words. Stars: binaries - supergiants - infrared: stars - X-rays: binaries

1. Introduction

The *INTEGRAL* satellite has broadened our knowledge of X-ray binaries, by discovering a significant population of new X-ray sources. The Third *IBIS/ISGRI* catalogue (Bird et al. 2007) contains more than 420 sources of which 167 are new. The on-line catalogue of *INTEGRAL* sources¹ lists 245 sources discovered or re-discovered by *INTEGRAL*. Of these, 61 ($\sim 25\%$) remain unidentified and 81 ($\sim 33\%$) are extragalactic objects (AGNs, Seyferts, QSOs, etc). The rest are mainly X-ray binaries. 51 of them ($\sim 21\%$) turn out to be High Mass X-ray Binary candidates (HMXB), which were missed by previous missions because of their high obscuration, because of their very short X-ray cycles or a combination of both. Approximately half of them have been confirmed so far. Amongst these new 51 HMXB systems, there are a few Be/X-ray binaries (BeXB), Supergiant X-ray binaries (SGXB) and the newly established class of Supergiant Fast X-ray Transients (SFXT; e.g., Smith et al. 2006; Negueruela et al. 2006a; Sguera et al. 2006). This last class comprises 16 candidates of which nearly half have been confirmed. Before we can

conclusively classify every new source and understand its X-ray behavior, we must first establish the nature of the counterpart.

Since the majority of these sources are located in the galactic plane, and concentrated towards the galactic center region, they suffer from large extinction ($A_V \lesssim 15 \text{ mag}$). Thus, their detection in the visual is often very difficult, and other strategies must be adopted. Detection and study of the counterparts in the infrared, however, is possible with 4 m class telescopes. An important caveat, though, is that the spectral classification of hot stars using infrared spectra is much more uncertain than using the well established MK standard 3950–4750 Å region. For example, the spectral classification based on the *K* band spectrum cannot be done without fundamental ambiguities because of the lack of adequate spectral features in that range (Hanson et al. 1996). This problem can be, at least partially, circumvented by using the combined information of several spectral bands. However, this is often impossible to achieve within a single observing campaign.

Following the discovery by *INTEGRAL*, intense observing campaigns have been undertaken (Chaty et al. 2008; Masetti et al. 2008; Nespoli et al. 2008; Negueruela et al. 2009, amongst others). As these discoveries proceed, it is fundamental to refine and establish the nature of these new sources. In

Send offprint requests to: J.M. Torrejón

¹ <http://isdcu13.unige.ch/~rodrigue/html/igrsources.html>

particular, a very important parameter is the distance to the system which, in turn, gives the X-ray luminosity, the main observable used to constrain the theoretical models. Most of the work done, however, has been based on *H* and/or *K* band spectra to overcome the high extinction. Classification based on a single infrared spectrum (in many cases of intermediate or low resolution only) allows only a crude approximation. For example, the difference in M_V between a B1 Ib and a B1 Ia supergiant (which are difficult or even impossible to tell apart based on an intermediate resolution *H* or *K* spectrum) results in an uncertainty of an order of magnitude error in L_X (Negueruela et al. 2009). In this paper, we combine new spectroscopy in several bands with new *JHK* photometry (also *RI* for one source) to further refine the spectral classification of a sample of these sources.

2. The observations

2.1. Photometry

JHK' band photometry was obtained with the 3.5 m Telescopio Nazionale Galileo (TNG) at the Island of La Palma (Canary Islands, Spain) during the nights of 2006 July 7 and 2007 May 1. In both runs the configuration was the same. The instrument used was the Near Infrared Camera and Spectrograph (NICS) in the Large Field configuration, equipped with a Rockwell 1024×1024 HgCdTe Hawaii array detector, which gives a spatial scale of 0''.25/pixel and covers a field of view of 4'.2×4'.2. During the night of 2006 July 7, the seeing was 0''.7 in the *J* band, very stable during the whole night, while on 2007 May 1, the seeing was 0''.8.

Each observation consists of a mosaic of five images constructed from a dithering pattern which kept the frame always on-source, using an automatic script available at the telescope. From this mosaic, flat-field correction, bad pixel map, sky subtraction and image co-addition were performed with the SNAP software². The photometry was subsequently performed using Starlink GAIA on the final co-added frame.

The Arnica standard star fields AS33 and AS27 (Hunt et al. 1998) were observed throughout the night at a range of airmasses. Transformation equations were applied to the program star giving the magnitudes listed in Table 1.

Comparison with previously published values show some discrepancies which can be readily explained. In particular, for IGR J19140+0951, comparison with 2MASS magnitudes shows a discrepancy. This is due to the close proximity of the bright star 2MASS J19149417+0952538, located very close to the south-east, which contaminates the 2MASS photometry (cf. in't Zand et al. 2006b). Any photometry made with a 3.5 m telescope under good seeing conditions must certainly be contaminated as well. However, careful attention has been paid here to the extraction to minimize this effect, making the magnitudes listed above rather more reliable than those of the 2MASS catalogue.

For IGR J19140+0951, *RI* band photometry was obtained with the Andalucia Faint Object Spectrograph and Camera (ALFOSC) on the 2.6 m Nordic Optical Telescope (NOT) in La Palma, on the night of 2006 June 18. The instrument was equipped with the thinned 2048×2048 pixel E2V CCD, covering a field of view of 6'.4 × 6'.4 with a spatial scale 0.19 arcsec/pixel. Standard Bessel *R* and interference *i* filters were used to observe both the target and the standard star fields. We obtain $R = 20.77 \pm 0.05$ and $I = 17.30 \pm 0.03$.

2.2. Spectroscopy

Infrared spectroscopy of several sources was performed with NICS. Medium resolution infrared spectroscopy in the *K* band was also obtained on 2006 July 7. The instrument was equipped with the K_b grism, which covers the 1.95–2.34 μm spectral region, and gives a resolution of 4.3 Å/pixel. A slit of 1'' was chosen, providing a resolving power $R = 1250$.

To estimate the sky, we observed the target stars integrating on four positions along the slit. We used the ABBA script which automatically moves the telescope, nodding the source along the slit, thereby producing a set of four images with typical exposure times ranging from 100 to 300 s for the programme stars thereby minimising the sky variability throughout each observation. The four images were median averaged and this median, subtracted from each individual image of the set. This method, together with the use of the 1'' slit, also minimizes any possible nebular contamination. These sky subtracted images were flat fielded and the resulting spectra averaged. The wavelength calibration was done with the Ar lamps available at the telescope. Analysis was subsequently performed with Starlink FIGARO and DIPSO.

A number of A0 V and G2 V stars were selected from the list available at the telescope and observed throughout the night at airmasses and positions as close as possible to the target fields.

For the telluric subtraction two methods were used. The first one, described in Hanson et al. (1996), makes use of the fact that spectra of A0 V stars are featureless in the *K* band except for the $\text{Br}\gamma$ line. A spectrum from the Sun was retrieved from the ESO archives. This spectrum was rebinned to match the sampling of our target spectra, wavelength shifted for the different radial velocities and, finally, the lines rotationally broadened to match the profiles of our G2 V standard stars. The spectrum of our G2 V stars was then divided by this corrected spectrum of the Sun. The resulting spectrum was subsequently used to correct the $\text{Br}\gamma$ zone in the spectrum of the A0 V standard star, isolating the true $\text{Br}\gamma$ profile of the early type star. Then, we fitted this line only and removed it from the A0 V spectrum, leaving a pure telluric spectrum which was subsequently applied to the programme stars.

The second method is described in Maiolino et al. (1996). Essentially, it uses directly the G2 V star spectrum divided by the modified solar spectrum to correct the whole spectrum of the program star. At the resolution of our spectra, these two methods give almost indistinguishable results. Where appropriate, we averaged the final spectra, of each source, obtained by both methods. The resulting spectra have been normalised to unity to allow for comparison with standard atlases.

In addition, infrared spectroscopy of the counterpart to IGR J19140+0951 was obtained using SPEX on the Infrared Telescope Facility (IRTF) on Mauna Kea on 2005 September 3. The *J*, *H* and *K* band spectra presented in Fig. 4 were constructed from medians of six 10 s exposures. The observing and data-reduction procedure are fully described in Harrison et al. (2004). SPEX was used in single-order mode with a 0''.3 slit, giving a dispersion of 5.51 Å/pixel. Unfortunately, the conditions at the IRTF were not photometric, but the seeing was excellent.

We employed a script with which data at six separate positions along the slit were obtained. To remove the sky background and dark current from each SPEX exposure, we subtracted the median of the other five exposures obtained in an observing sequence. This process resulted in six background-subtracted exposures from which the spectra were extracted using the normal IRAF methods. The spectra were wavelength calibrated by the

² <http://www.tng.iac.es/news/2002/09/10/snap/index.html>

Table 1. *JHK'* infrared photometry for the sources included in our survey. The corresponding 2MASS counterparts and the reference of their discovery are also included.

Source	2MASS	R.A.	Dec.	Ref.	<i>J</i> (1.27 μ m)	<i>H</i> (1.63 μ m)	<i>K'</i> (2.12 μ m)
IGR J18027–2016	2MASS J18024194–2017172	18 02 41.94	–20 17 17.3	C	12.79 $\pm$ 0.06	11.96 $\pm$ 0.08	11.48 $\pm$ 0.05
SAX J18186–1703	2MASS J18183790–1702479	18 18 37.90	–17 02 47.9	Z1	10.08 $\pm$ 0.06	8.94 $\pm$ 0.08	7.79 $\pm$ 0.05
IGR J18483–0311 ^a	2MASS J18481720–0310168	18 48 17.20	–03 10 16.8	C	10.84 $\pm$ 0.03	9.38 $\pm$ 0.02	8.47 $\pm$ 0.02
IGR J19140+0951	2MASS J19140422+0952577	19 14 04.23	+09 52 57.7	Z2	11.33 $\pm$ 0.06	9.71 $\pm$ 0.08	8.83 $\pm$ 0.05

C: Chaty et al. (2008)

Z1: in't Zand et al. (2006a)

Z2: in't Zand et al. (2006b)

^a Photometry from Rahoui & Chaty (2008). For this source, the filters are centered at 1.25, 1.65 and 2.20 μ m respectively.

extraction of an Ar arc spectrum at the position (aperture) of each spectrum. We observed the G2 V star HD 150698 (in an identical fashion) to correct for telluric features as described above.

Finally, *I*-band spectra of IGR J18027–2016 and IGR J18483–0311 were obtained on 2008 May 16 using the 4.2-m William Herschel Telescope (WHT), in La Palma (Spain) equipped with the ISIS double-beam spectrograph, during a service run. We used the red arm fitted with the R316R grating and the RED+ CCD, a configuration that results in a nominal dispersion of 0.85 Å/pixel (the resolution element is approximately 3 pixels). We also observed the counterpart to IGR J19140+0951 on 2008 May 18 using the same instrument, but this time fitted with the R158R grating, which results in a nominal dispersion of 1.8 Å/pixel.

3. Results

For the line identification and spectral classification we have used the following atlases: Andriolat et al. (1995) for the *I* band, Wallace et al. (2000) for the *J* band; Blum et al. (1997), Meyer et al. (1998) and Hanson et al. (1998) for the *H* band; Hanson et al. (1996) and Hanson et al. (2005) for the *K* band.

As has been pointed out, the classification of massive stars using the 2 micron spectral range is difficult due to the lack of spectral features in this domain. In the case of HMXB the situation is still worse because the spectra of the infrared counterparts might be altered by the presence of the compact object in the system. Little work has been done in the near-infrared observations of HMXBs (see, for example, Clark et al. (1999); Fig. 7, for some BeXs). In the pioneering work by Hanson et al. (1996) a number of HMXB are included (cf. Fig. 29 in that reference). In almost all of them, the relevant diagnostic lines appear in emission, both in BeXs and in SGXBs. However, isolated supergiants tend to show these lines in absorption (Hanson et al. 1996). No HMXBs are included in the higher S/N survey of Hanson et al. (2005). Notwithstanding, this last work has been used already to classify the infrared counterparts of some HMXBs (Mason et al. 2009, *vg.*). Lacking a comprehensive scheme to classify HMXB counterparts from near-infrared spectra, we will use the previously cited available atlases. We estimate that the uncertainty due to the above mentioned effects alone could amount to half a spectral subtype.

In Table 2 we give the equivalent widths of the main diagnostic lines found in our spectra. Due caution must be exercised, however, in using them for spectral classification purposes, as stated by Hanson et al. (2005, ;see their Section 5.2) as only the comparison between different classes appears to be meaningful.

In order to compute their distance and L_X we will use the infrared distance modulus computed from the following equation.

$$K_0 - M_K = 5 \log d - 5, \quad (1)$$

where $K_0 = K - A_K$ are the infrared magnitudes corrected for extinction and K magnitudes are taken from Table 1. We compute the M_K band absolute magnitudes from the $(V - K)_0$ colors given by Ducati et al. (2001) taking into account the M_V values from Schmidt-Kaler (1982). To compute the total to selective absorption A we will first compute the corresponding IR color excess $E(J - K) = (J - K) - (J - K)_0$, where the intrinsic IR colors are taken from Ducati et al. (2001) for the deduced spectral type. These excesses are converted to $E(B - V)$ by means of the following relation $E(J - K) = 0.50E(B - V)$. Finally $A_K = 0.36E(B - V)$ (Fitzpatrick 1999), assuming the standard extinction law ($R_V = 3.1$). We compute the distance for the two extremes of the range in spectral type compatible with our observations, namely, the cold-faint and hot-luminous ends. The range of possible distances so deduced and the corresponding X-ray luminosities are given in Table 3. The quoted errors refer only to the dispersion of these estimated values. The final uncertainties in the distance will be dominated by the intrinsic dispersion of the calibration of the order of 0.5 mag.

3.1. IGR J18027–2016

This source was discovered by Revnivtsev et al. (2004) using *INTEGRAL*. The source has a very large hydrogen column ($N_H = (6.8 \pm 1) \times 10^{22} \text{ cm}^{-2}$). It has been identified with SAX J1802.7–2017, which was discovered serendipitously by Augello et al. (2003) with *BeppoSAX*. Using *XMM-Newton*, Walter et al. (2006) localised the source position at $\alpha = 18 \text{ h } 02 \text{ m } 42.0 \text{ s}$, $\delta = -20^\circ 17' 18''$ with a 4'' positional accuracy. Its 2–10 keV unabsorbed flux during the primary pulse is $8.9 \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$. It is an eclipsing X-ray pulsar ($P_{\text{spin}} = 139.612 \text{ s}$) which revolves the donor star each 4.6 d (Hill et al. 2005).

Based on the orbital solution, Hill et al. (2005) proposed that the mass donor should be an early type supergiant with most likely parameters $M_* \sim 21 M_\odot$ and $R_* \sim 19 R_\odot$. Masetti et al. (2008) identified the counterpart as 2MASS J18024194–2017172 and obtained a low-resolution optical spectrum that showed it to be a very reddened early-type star. Using the mass and radius deduced by Hill et al. (2005), they argue that the counterpart should be a B-type giant. Chaty et al. (2008) confirmed the identity of the counterpart.

Table 2. Equivalent widths ($\text{\AA}$) of the main diagnostic lines in the *JHK* bands observed in our sample. Typical errors are always less 1 $\text{\AA}$. Numbers beside chemical species are the laboratory wavelengths in μm . *I* band equivalent widths have not been measured due to the line blending.

Source	He I 2.058 μm	He I 2.1126	N III 2.1155	Br γ 2.1661	He I 1.083	Pa β 1.2818	Br 11 1.6814	He I 1.7004
IGR J18027–2016	4.1(a+e) -4.5(e)	2.6	-0.2	1.1 (a+e) -4.1 (e)	...	...	...	...
SAX J18186–1703	8.6(a+e) -6.5(e)	2.0	...	5.1	...	...	...	...
IGR J19140+0951 [†]	-1.9(a+e) -5.7(e)	2.2	...	2.0(a+e) -1.0(e)	...	...	...	...
IGR J19140+0951 [‡]	-1.1(a+e) -1.3(e)	2.2	-0.4	1.2(a+e) -2.6(e)	-7.1	1.8(a+e) -5.3(e)	2.3	2.1

(a+e): Absorption + Emission; (e): Emission only

[†] TNG data; [‡] SPEX-IRTF data




Fig. 1. **Left:** *WHT I*-band spectrum of IGR J18027–2016 normalised to flux unity bracketted between comparison standard stars. **Right:** *TNG K*-band spectrum normalised to flux unity. Emission is clearly seen in Br γ and He I lines, and also likely in N III 2.1155 μm , suggesting an early B supergiant.

Their infrared spectrum seems typical of an early-B supergiant, displaying prominent Brackett lines in the *H* band even though these authors claimed the detection of He I and He II lines in the optical spectrum. In fact, they favor a B type supergiant based on a $T_{\text{eff}} \sim 20800$ K deduced from a fit to the SED. Based on the R_*/D value of the normalization, and assuming a stellar radius $R_* = 20 R_{\odot}$, they estimate a distance of ~ 12 kpc. Chaty et al. (2008) estimate $A_V \sim 8.8$ mag, while Masetti et al. (2008) estimate ~ 8.3 mag. In the following, we will use our TNG *K*-band

and *WHT I*-band spectra (Fig. 1) to refine the classification of the counterpart and establish the nature of the system.

K BAND SPECTRUM. The presence of He I lines and the absence of He II (which is seen up to O9) points towards a B type star. The Br γ line is in emission. This line seems to be blended with the blueward absorption of the He I 2.161 μm line. This morphology is more pronounced in high luminosity B stars (Hanson et al. 2005, Fig. 12). He I 2.058 μm is also in emission, with a possible P-Cygni profile, a feature also noted by Chaty et al. (2008) on their lower resolution spectrum. In the atlas of Hanson et al.

(2005), no luminosity class III star shows He I 2.058 μm in emission. In contrast, this line is in emission in B supergiants and Be stars. However, He I 2.113 μm is not seen in Be stars, while it is very prominent in our object, a feature of B supergiants (see Figures 13 to 16 in Hanson et al. 2005). We therefore conclude that the counterpart is an early B supergiant star.

I BAND SPECTRUM. As seen in Fig. 1, our *I*-band spectrum has a low SNR. However, it is easy to see the narrow well-defined Paschen lines, typical of a high-luminosity object. Moreover, Paschen lines are visible up to Pa 19, in spite of moderate SNR and resolution, clearly identifying the star as a supergiant (compare to the B1 V star in the same figure). The prominent He I lines are typical of supergiants with spectral types close to B2 (compare to the earlier supergiants in the same figure). Moreover, as can be seen in the spectra of the O9.5 Ia, for stars B0.5 and earlier, the C III 8502 Å is blended with Pa 16, making it stronger than the neighbouring Pa 15 and Pa 17. This is not the case in our object, suggesting it is later than B0.5. On the other hand, the red spectrum of IGR J18027–2017 does not display O I 7774 Å (which is not seen in the spectrum of Chaty et al. (2008) either). This would indicate a type earlier than B1.5 if it is a luminous supergiant or earlier than B2 if it is less luminous.

Taking together all the data, we propose a B1 Iab - B1 Ib supergiant counterpart. To discriminate between the two luminosities is very difficult with the available spectra. However, the eclipse duration covers ~ 0.25 of the orbit (Hill et al. 2005). With a period of only 4.6 d, a B1 Iab supergiant would not fit into the orbit. Therefore, a luminosity class Ib is definitely assigned to the donor. This locates the source at a distance of $d \approx 12.4$ kpc, in agreement with Chaty et al. (2008), between the Sagittarius and Perseus arms, well beyond the Galactic Center (Fig. 8).

Given this classification, the observed color excess is $E(B - V) \approx 3$ and, assuming a standard extinction law, the extinction in the visual band would be $A_V \approx 9.4$. This is slightly larger than the value given by Chaty et al. (2008) ($A_V \approx 8.8$) and by Masetti et al. (2008), namely $A_V \approx 8$.

For a standard reddening law (Fitzpatrick 1999), and typical intrinsic colours $(I - J)_0 = -0.15$, $(R - I)_0 = -0.17$ (Ducati et al. 2001), our value $E(J - K) \approx 1.54$ implies $I \approx 14.8$ and $R \approx 16.9$ in excellent agreement with the value $R = 16.9 \pm 0.1$ measured by Masetti et al. (2008).

3.2. SAX J18186–1703

SAX J1818.61703 was discovered by *BeppoSAX* during a strong outburst with a fast rise time of ~ 1 h, in March 1998 (in’t Zand et al. 1998). Subsequently *INTEGRAL* detected a double-peaked outburst in September 2003 (Grebenev & Sunyaev 2005) and two more in October 2003 (Sguera et al. 2005). Other fast outbursts have been observed with the ASM onboard *RossixTE* (Sguera et al. 2005). Its counterpart, 2MASS J18183790–1702479, was identified by a *Chandra* localization (in’t Zand et al. 2006a). Recently Zurita-Heras & Chaty (2009), using *INTEGRAL*, have reported a 22–50 keV flux of the order of $(2-8) \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$ in quiescence while it reaches $(1-2) \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$ in the strongest flares. No pulsations have been detected so far, but an orbital period of 30 d has been established (Zurita-Heras & Chaty 2009).

K BAND SPECTRUM. In Fig. 2, we present the *K* band spectrum of the IR counterpart. The absence of He II 2.1885 μm confirms that this star is later than $\sim \text{O8}$. The He I 2.0581 μm line




Fig. 2. *K* band spectrum of the counterpart to SAX J18186–1703 with the most relevant features expected in this range identified. Note the absence of any feature at the position of the He II 2.1885 μm line.

is strongly in emission. This is typical of early B supergiants. Along with the morphology of the complex He I 2.1607/2.1617 - Bry it points to high luminosity. Main sequence early B stars have broader lines (Hanson et al. 2005, Fig. 12). Following the same arguments presented for IGR J18027–2017, we assign a similar spectral type to this object and conclude it is a supergiant in the B0–B1.5 range. The exact luminosity class is difficult to determine, but, as discussed previously, it is likely to be at least Iab. This would place the source at a distance of $d \approx 2.1$ kpc in the outskirts of the Sagittarius arm (Fig. 8).

3.3. IGR J18483–0311

IGR J18483–0311 is a transient X-ray source discovered by Chernyakova et al. (2003) during observations of the Galactic Center with *INTEGRAL*. Using *Swift*, Sguera et al. (2007) suggest as a counterpart the IR source 2MASS J18481720–0310168, which is confirmed by Chaty et al. (2008). The X-ray source shows very high obscuration ($N_H = 9^{+5}_{-4} \times 10^{22} \text{ cm}^{-2}$). The system contains a pulsar ($P_s = 21.0526$ s) in orbit around the primary with a period of $P_{\text{orb}} = 18.52$ d (Sguera et al. 2007). The unabsorbed X-ray 20–100 keV flux, measured with IBIS on board *INTEGRAL*, is $\sim 2 \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$ (Sguera et al. 2007).

By comparing a low-resolution *HK* spectrum to spectra of classification standards, Rahoui & Chaty (2008) conclude that the counterpart is a B0.5 supergiant, and suggest a luminosity class Ia. In Fig. 3, we present the *I* band spectrum of this object, bracketed between those of two supergiants with well-determined spectral type. Our spectrum shows narrow well-defined Paschen lines, typical of a high-luminosity object. Moreover, Paschen lines are visible up to Pa 18, in spite of moderate SNR and resolution, clearly identifying the star as a supergiant (compare to the B1 V star in the same figure). The pres-




Fig. 3. *I* band spectrum of IGR J18483–0311, compared to several other objects with well-determined spectral types. Note the similarity of the spectrum to the B1 Ib star. Note also the strong DIBs, indicating a heavy interstellar absorption.

ence of strong He I 8779 Å is typical of early B supergiants, while the absence of O I 8446 Å and N I lines makes it earlier than B2. Pa 16 is not prominently stronger than Pa 17, suggesting that it does not contain an important contribution from C III 8502 Å. As discussed above, this suggests a spectral type B0.5 or later. The exact determination of the luminosity class is not easy with the data available. However, the fact that we only detect Paschen lines up to P18 suggests that the star is not a Ia supergiant. Combining our data with the *HK* spectrum of Rahoui & Chaty (2008), we conclude that the star is a B0.5–B1 supergiant, most likely with a luminosity class Iab. This would locate the system at $d \sim 2.8$ kpc.

3.4. IGR J1914.0+0951

IGR J19140+0951 was discovered by Hannikainen et al. (2004) with *INTEGRAL*. It displays X-ray characteristics typical of a SGXB. in’t Zand et al. (2006b) used *Chandra* to pinpoint the IR counterpart, which turns out to be 2MASS J19140422+0952577. A period of 13.55 d is observed with *RossixTE* (Corbet et al. 2004). No pulsations have been detected so far.

in’t Zand et al. (2006b) presented a low resolution red spectrum of the source, that was very badly affected by fringing. It shows no counts bluewards of 7000 Å. They also quote a magnitude $I = 13.0$ for the counterpart, completely at odds with our value, and the low counts in the corresponding region of the spectrum.

Unfortunately, our *I*-band spectrum, though not affected by fringing is of low resolution and SNR. We identify a few narrow, strong Paschen lines, that indicate an early-type supergiant. We

do not detect the O I 7774 Å line, which (given the low SNR in this part of the spectrum) constraints it to be earlier than ~B3.

Hannikainen et al. (2007) have presented a low SNR *K*-band spectrum and an *H*-band spectrum of much better resolution and SNR. Their analysis led them to conclude that the star was a B0.5 Ia supergiant. Nespoli et al. (2008) classify it as B1 Iab, based on a *K*-band spectrum. The spectrum from Hannikainen et al. (2007) displays Br γ weakly in absorption, while this line is more clearly in absorption in the spectrum of Nespoli et al. (2008). However, our TNG spectra, taken four months later, as well as our SPEX spectra, taken one year earlier (both of higher SNR), show a P-Cygni profile in this line. This is indication of high activity in this object and further supports the high luminosity of the star.

No He II absorption lines are seen in our spectra, suggesting that the star is later than O9. The presence of a very well defined series of Brackett H I lines in the *H* band (see Fig. 4, *H* band panel) also points towards an early B star, as these lines disappear for O type stars. The narrowness and depth of the Br lines is a clear indication of high luminosity. Moreover, we detect them well up to Br19, at least. The narrow and deep He I 1.7 μ m line is also typical of a supergiant. It is deeper than neighbouring Br lines, indicating high luminosity. Its relative depth with respect nearby Br 11 points clearly towards type B0–B0.5 Ia and rules out types earlier than O9.7I (Hanson et al. 1998) (Fig. 2).

In addition, we find He I 2.0581 μ m strongly in emission, a characteristic of luminous early-B supergiants. This emission seems to be variable, as it is much more pronounced in the TNG data than in the IRTF observations (Fig. 4). We also see emission in the He I 1.083 μ m line in the *J* band, and in Pa β .

We conclude that the spectral type is B0.5 Ia, compatible with that given by Hannikainen et al. (2004) and Nespoli et al. (2008), but with a slightly higher luminosity class. Nespoli et al. (2008) estimate a distance of ≈ 1.1 kpc, but they use the 2MASS photometry, contaminated by the very bright star towards the upper right (see Fig. 5), namely 2MASS J19140417+0952538 ($K_s \approx 6.5$ mag), as was already recognised by in’t Zand et al. (2006b). We performed our photometry on images of higher spatial resolution which allowed us to clearly resolve both stars. We used an aperture small enough to ensure that the contribution from the bright star is minimal. Then we applied aperture correction to our magnitudes keeping in mind that avoiding completely the contamination is probably impossible without adaptive optics. Using our photometry, rather than the 2MASS values, and the deduced spectral type, we compute a distance of $d \approx 3.6$ kpc.

In Fig. 6 we make use of our red and infrared photometry for this object to plot the *RIJHK* photometry absolute flux calibration, dereddened with a standard reddening law and taking $E(B - V) = 5.5$. The photometry is superimposed on a Kurucz atmosphere model for $T_{\text{eff}} = 28000$ K and $\log g = 3.5$, corresponding to an early type B supergiant³. The acceptable fit shows that the extinction to this source is close to standard, in spite of the high value. The normalization constant in Fig. 6, assuming $R_* \approx 30 R_\odot$, would yield a distance of ≈ 3.6 kpc, in agreement with the value computed previously. The independent estimate by Rahoui & Chaty (2008), based on a fit to the near- and mid-infrared magnitudes, of $d \sim 3.1$ kpc, is also consistent with our value.

³ Actually, $\log g = 3.0$ would be more adequate. Unfortunately, this Kurucz model is not available. The difference in slope, however, is negligible for the present purposes




Fig. 4. SPEX-IRTF *J*, *H* and *K* band spectra of IGR J19140+0951. In the *K* band, TNG data (top) are also presented.




Fig. 5. The counterpart to IGR J19140+0951. **Left panel:** *K*-band image from 2MASS, showing the contamination by the light of the bright star 2MASS J19140417+0952538, which affects the catalogue photometry for this source. **Right panel:** TNG *K'* band image, showing the two stars clearly resolved (cf. Fig. 2 in Rahoui & Chaty (2008).)

3.5. XTE J1901+014

The source XTE J1901+014 was discovered by the *RXTE*/ASM monitor in 2002 during a brief outburst, after which the source went into quiescence below detectable limits. Examination of archival data showed that the source had undergone a previous brief outburst in 1997 (Remillard & Smith 2002) showing a behaviour reminiscent of that of SFXTs. Smith et al. (2007) used *XMM-Newton* to obtain accurate position of the X-ray source, which turns out to be $\alpha = 19^{\text{h}} 01^{\text{m}} 40.20^{\text{s}}$, $\delta = +01^{\circ} 26' 26''.0$ with an error of $\sim 1''$. Subsequently we searched the area for possible counterparts. In Fig. 7, we show a *K'*-band image of the area encompassing the error circle. Only one possible candidate is visible within the *XMM-Newton* error circle. The object is very faint with $K' \gtrsim 14.2$, close to the limit of the sky brightness at $2\mu\text{m}$ at the La Palma Observatory. No sources show up in the *J*- or *H*-band images.

As the infrared observations do not provide enough information, in order to guess the nature of the system we will resort to X-ray data. The *XMM-Newton* spectra can be well fitted by a thermal Bremsstrahlung with $kT \sim 6$ keV and an absorption column of $N_{\text{H}} \sim 2.5 \times 10^{22} \text{ cm}^{-2}$ (Rampy 2009, priv. comm.). This would correspond to $E(B - V) \approx 3.7$ (Ryter 1996), which is larger than the integrated value for the Milky Way Galaxy in the direction of the source, namely $\sim 8.5 \times 10^{21} \text{ cm}^{-2}$ suggest-

ing intrinsic absorption, as might be expected. Assuming that all absorption is interstellar, the corresponding infrared excess is $E(J - K) \approx 1.8$. Comparing these values with those shown in Table 3, we can see that the source is, indeed, very reddened but not particularly so.

If we assume an M_K typical for a not very luminous OB supergiant (~ -5.5 ; Martins & Plez 2006), with $K \approx 14.2$, the distance to the source would be ~ 47 kpc, placing it well outside the Galaxy. This seems very unlikely. At the outskirts of the Galaxy in the direction to the source (~ 22 kpc), the infrared magnitude of the counterpart would be $M_K \approx -3.8$. This would correspond to a O8 V star or a K5 III star. Therefore the possibility of a supergiant companion is ruled out.

However, there are arguments to believe that the distance must be much lower. Indeed, the 2002 outburst detected with *RossixTE*/ASM reached a luminosity of the order of 1 Crab, which corresponds to a flux $F_X = 3.63 \times 10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$. At a distance $d = 22$ kpc, the outburst should have had an $L_X \approx 2 \times 10^{39} \text{ erg s}^{-1}$. This is equivalent to the Eddington luminosity for a compact object of $\sim 16 M_{\odot}$, and higher than ever reported for any neutron star system. On the other hand, if we assume that the outburst luminosity was at the Eddington luminosity for accretion onto a NS ($\sim 2 \times 10^{38} \text{ erg s}^{-1}$), the maximum distance to the source should be of the order of $d_{\text{max}} \sim 7$ kpc, which would place the source at the tip of the Long Bar in the




Fig. 6. Our *RIJHK* photometry of IGR J19140+0951 dereddened with a standard extinction law and $E(B-V) = 5.5$ appears superimposed on a Kurucz atmosphere model for $T_{\text{eff}} = 28000$ K and $\log g = 3.5$, corresponding to an early type B supergiant.

galactic nucleus. For such a source, $M_K \approx -1.28$ which would be compatible with a $\sim$ G5 III star. In fact, the maximum distance at which the primary would be compatible with a class III star is $d \sim 5$ kpc, in which case, we would have G4III (or also B4V). Below this distance, essentially, only main sequence stars are allowed. For distances around 1 kpc or less, the only compatible companions would be G, K or M main sequence stars. This raises the interesting possibility that the system is, actually, a Cataclysmic Variable, since the X-ray spectrum can be described by a thermal Bremsstrahlung at 6 keV. However, in such a case, the $L_X \sim 10^{36} \text{ erg s}^{-1}$ which would be several orders of magnitude larger than that usually found in CV. Therefore, we conclude that, very likely, $d_{\text{min}} \sim 1 - 2$ kpc. This is in good agreement with the conclusions obtained by Karasev et al. (2008), who argue in favor of a distance of ~ 5 kpc.

4. Discussion and conclusions

Four of our sources have early-B supergiant companions. All four counterparts turn out to cover a rather narrow spectral interval, being early-B (B0–B1.5) supergiants of moderate or high luminosity. This identifies the nature of the X-ray source unambiguously as HMXBs. Furthermore, they are all heavily obscured, with $E(B-V) \sim 3-5.5$, implying extinctions on the order of $A_V = 9 - 16$ mag in the visual band. But, on the other hand, XTE 1901+014 has no obvious counterpart in any band except K' . Thus, a supergiant nature is definitely excluded, thereby setting it apart from the other four.

IGR J18027–2017 is a persistent X-ray source with a counterpart around B1 Ib. It is therefore, a SGXB system. The reason why it has not been detected by earlier X-ray missions is not clear. For example, it was not detected by *ROSAT*. At a distance of $d \approx 12.4$ kpc, its *XMM-Newton* detected flux in the 2–10 keV band for the main pulse of $8.9 \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$, would yield




Fig. 7. TNG K' -band image of the field around XTE J1901+014. The blue circle represents the nominal *XMM-Newton* position with an error radius of $1''$ (which for our image scale is 4 pixels). A faint possible candidate is seen inside the *XMM-Newton* error circle. The candidate must have $K' \gtrsim 14.2$. It is not visible in either J or H bands.




Fig. 8. Sketch of the Galaxy showing the position of the HMXBs studied in this work (filled dots). Numbers refer to the relative order of the source in Table 3. Galaxy model from Churchwell et al. (2009).

a luminosity of $L_X = 1.65 \times 10^{36} \text{ erg s}^{-1}$. This is typical for these kind of systems albeit in the lower end. Since the detected flux outside the primary pulse is a little bit smaller, the system can spend a fraction of its time in the upper $\times 10^{35} \text{ erg s}^{-1}$. Given its high obscuration, this can explain why it was missed by earlier surveys.

For SAX J18186–1703, the NIR counterpart is a B0.5Iab star, confirming its nature as a SFXT. The 22–50 keV flux reported by Zurita-Heras & Chaty (2009) is of the order of $(2-8) \times 10^{-11} \text{ erg s}^{-1} \text{ cm}^{-2}$ in quiescence while it reaches $(1-2) \times 10^{-9} \text{ erg s}^{-1} \text{ cm}^{-2}$ at the strongest flares. At the deduced dis-

Table 3. Derived parameters for stars in our sample, calculated using the observed data.

Source	Spectral type	Class	$E(B - V)$	$E(J - K)$	M_V	M_K	d (kpc)	L_X (erg/s) ^(a)
IGR J18027–2016	B1 Ib	SGXB	3.04±0.02	1.54	−5.8	−5.1	12.4±0.1	$(1.65±0.03) \times 10^{36}$
SAX J18186–1703	B0.5 Iab	SFXT	5.08±0.05	2.54	−6.4	−5.65	2.1±0.1	$(8±0.4) \times 10^{35}$
IGR J18483–0311	B0–1 Iab	SFXT	5.22±0.02	2.61	−6.4	−5.66	2.83±0.05	$(1.9±0.06) \times 10^{36}$
IGR J19140+0951	B0.5 Iab-a	SGXB	5.5±0.1	2.75	−6.65±0.03	−5.90±0.25	3.6±0.04	$(1.6±0.3) \times 10^{36}$
XTE 1901+014	G5 III - A3 V	LMXB?	~3.7	1.8			2-7	$1^{+1}_{-0.8} \times 10^{38}$

^a Observed peak luminosities for the energy ranges (2–10) keV, (22–55) keV, (20–100) keV, (2–20) keV and (3–12) keV respectively.

tance of ~ 2.1 kpc, the X-ray luminosity of this object would be $\sim 3 \times 10^{34}$ erg s^{−1} in quiescence, while the peak luminosity can be as high as $\sim 8 \times 10^{35}$ erg s^{−1}. This is lower than those found in other SFXTs ($\sim 10^{36}$ erg s^{−1}). Zurita-Heras & Chaty (2009) have found that this system is in a very eccentric orbit ($e \sim 0.3 - 0.4$), with $P_{orb} \approx 30$ d, amongst the largest yet found for SFXT, reaching a periastron distance between 2 and 3 R_* . This would locate the compact object at a slightly larger distance from the donor than in the SGXB systems ($a \lesssim 2R_*$) thereby reducing the L_X of the outbursts slightly below 10^{36} erg s^{−1} as is observed.

IGR J18483–0311 is a transient system with a B0–B1 Iab primary. It has been classified as an intermediate SFXT (Rahoui & Chaty 2008). Like many other SFXTs, it is a nearby source at ~ 2.8 kpc. For this distance, the *INTEGRAL*/IBIS 20–100 keV flux (Sguera et al. 2007) translates into a luminosity $\sim 1.9 \times 10^{36}$ erg s^{−1}, for the strongest flares, which is typical for this kind of objects. The orbital period of this system is 18.5 d, while the spectral type of the companion is similar to that of SAX J18186–1703. Therefore, the compact object would be located slightly closer to the star and, correspondingly, its average X-ray luminosity will be higher, in agreement with the scenario described in Negueruela et al. (2008).

IGR J19140+0951 is a persistent X-ray source and therefore, owing to its counterpart, a SGXB located at $d \approx 3 - 4$ kpc, a little bit further than previously thought. For the bright *INTEGRAL* flux in the 2-2-0 keV band of 1×10^{-9} erg s^{−1} cm^{−2} (Hannikainen et al. 2004), the $L_X = 1 - 2 \times 10^{36}$ erg s^{−1}, typical of a SGXB albeit, again, in the lower end. The fact that, together with IGR 18027–2017, they have lower X-ray fluxes than the average SGXBs explains their more recent detections in the newer, and deeper *INTEGRAL* observations.

As can be seen in Fig. 8, the sources tend to concentrate in or behind the section of the Sagittarius arm projected onto the direction of the Galactic Bulge, an area intensively scanned by *INTEGRAL*. They are, thus, relatively close to the Sun (in comparison to typical distances to HMXBs), although highly reddened. This fact can introduce a bias in the newly detected populations of obscured sources, which tend to harbor SG companions, as only the brighter and/or closer ones can be reached with telescopes of moderate apertures in the 4 m class. A transient system with a main sequence companion, like a BeXB, will be too faint to be conclusively classified (i.e. Zurita-Heras & Chaty 2008). Remarkably, the furthest source in our sample is the least reddened one. This raises the question of the true abundance of HMXBs (and OB stars in fact) in our Galaxy. As can be seen in Table 3, the high reddening is not a sufficient condition to be an SFXT since some SGXBs (like IGR J19140+0951) are more reddened. On the other hand, it is not a *necessary* condition either, as other SFXTs present rather low reddenings (for example

IGR J08408–4503 with $E(B - V) = 0.5$ Negueruela et al. 2009). Furthermore, the position in the arms of the Galaxy of the newly discovered HMXBs (including the SFXTs) is entirely consistent with the positions of the previously known HMXBs (including SGXBs). Therefore there is no reason to believe that they form a different population. This would be expected from a scenario like that discussed in Negueruela et al. (2008). In this scenario, the SFXTs are explained as just SGXBs where the compact object is located further out in the system, in a region where the stellar wind of the supergiant companion has become substantially clumped. The number of HMXBs in the Galaxy would then be much larger than previously thought. But so would the number of isolated OB stars whose much lower X-ray emission will not show up in the current X-ray surveys if they are heavily obscured. This is supported by the discovery, in the last few years, of a growing population of very massive infrared clusters (e.g. Crowther et al. 2006; Messineo et al. 2008). Therefore, in principle, there is no reason to believe that the relative abundance of HMXBs, with respect to OB stars, differs from the current values which are well explained by the theoretical models.

Finally, XTE J1901+014 emerges as a transient source whose X-ray behaviour is characteristic of the SFXTs but where the presence of a Supergiant companion is clearly ruled out. It can still be a massive star close to the main sequence in the outskirts of the Galaxy, perhaps with a $16 M_\odot$ BH as a companion. In this respect, it is interesting to note the case of the LMC transient A 0535–66 reaching $L_X \approx 8 \times 10^{38}$ erg s^{−1} (e.g., Pakull & Parmar 1981). A similar source at the edge of the Galaxy could have a similar hard X-ray behaviour. However, taking into account the moderate hardness (the source is seen as 1RXH J190140.1+012630) and obscuration of the source, it seems more likely that the counterpart is a late type star located between 2 kpc and the Long Galactic bar (7 kpc), somewhere between luminosity class V to class III, but definitely not consistent with a class I star. Whether a main sequence or giant late type star wind can produce the instabilities necessary to produce the observed outbursts via accretion onto a compact object remains an issue.

Acknowledgements. This research has been funded by grants AYA2008-06166-C03-03 and Consolider-GTC CSD-2006-00070 from the Spanish Ministerio de Ciencia e Innovación (MICINN). JMT also acknowledges the research grant PR2007-0176. The authors would like to thank the referee, Dr. Alex Kaper, for his useful comments and suggestions.

The Nordic Optical Telescope is operated on the island of La Palma jointly by Denmark, Finland, Iceland, Norway, and Sweden, in the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias. The data were taken with ALFOSC, which is owned by the Instituto de Astrofísica de Andalucía (IAA) and operated at the Nordic Optical Telescope under agreement between IAA and the NBIfAFG of the Astronomical Observatory of Copenhagen.

The WHT is operated on the island of La Palma by the Isaac Newton Group in the Spanish Observatorio del Roque de Los Muchachos of the Instituto de Astrofísica de Canarias. The WHT data presented here have been obtained as part of the service programme. We thank the support astronomers for their dedication during the service nights.

Partly based on observations made with the Italian Telescopio Nazionale Galileo (TNG) operated on the island of La Palma by the Fundación Galileo Galilei of the INAF (Istituto Nazionale di Astrofisica) at the Spanish Observatorio del Roque de los Muchachos of the Instituto de Astrofísica de Canarias.

This research has made use of the Simbad data base, operated at CDS, Strasbourg (France) and of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

References

- Andrillat Y., Jaschek C., Jaschek M. 1995, A&AS, 112, 475
- Augello, G., Iaria, R., Robba, N.R., et al. 2003, ApJ, 596, 63
- Bird A.J., Malizia A., Bazzano A. et al. 2007, ApJ 170, 175
- Blum, R.D., Raymond, T.M., Conti, P.S., Figer, D.F., & Sellgren, K. 1997, AJ, 113, 1855
- Chaty, S., Rahoui, F., Foellmi, C., et al. 2008, A&A, 484, 783
- Churchwell E., Babler B.L., Meade M.R. et al. 2009, PASP 121, 213
- Chernyakova, M., Lutovinov, A., & Capitanio, F. 2003, ATel 157
- Clark J.S., Steele I.A., Fender R.P., Coe M.J. 1999, A&A 348, 888
- Corbet, R.H.D., Hannikainen, D.C., & Remillard, R. 2004, ATel 269
- Crowther P.A., Hadfield L.J., Clark J.S., Negueruela I., Vacca W.D. 2006, MNRAS 372, 1407
- Ducati, J.R., Bevilacqua, C.M., Rembold, S.B., & Ribeiro, D. 2001, ApJ, 558, 309
- Fitzpatrick, E.L. 1999, PASP, 111, 63
- Grebenev, S.A., & Sunyaev, R.A. 2005, Ast. Lett., 31, 672
- Halpern, J.P., & Gotthelf, E.V., ATel 344
- Halpern, J.P., & Gotthelf, E. V. 2007 ApJ, 669, 579
- Hannikainen, D.C., Rodriguez, J., Vilhu, O. et al. 2005, A&A, 435, 995
- Hannikainen, D.C., Rawlings, M.G., Muhli, P., Vilhu, O., Schultz, J., & Rodriguez, J. 2007, MNRAS, 380, 665
- Hanson, M.M., Conti, P.S., & Rieke, M.J. 1996, ApJS, 107, 281
- Hanson, M.M., Rieke, G.H., Luhman, K.L. 1998, AJ 116, 191
- Hanson, M.M., Kudritzki, R.P., Kenworthy, M.A., Puls, J., & Tokunaga A.T. 2005, ApJS, 161, 154
- Harrison T.E., Howell S.B., Szkody, P., Homeier D., Johnson, J.J., Osborne, H.L. 2004, ApJ 614, 947
- Hill, A.B., Walter, R., Knigge, C., et al. 2005, A&A, 439, 255
- Hunt, L.K., Mannucci, F., Testi, L., et al., AJ, 115, 2594
- in't Zand, J., Heise, J., Smith, M., et al. 1998, IAU 6840, 2
- in't Zand, J.J.M., Jonker, P., Mendez, M., & Markwardt, C. 2006a, ATel 915
- in't Zand J.J.M., Jonker P.G., Nelemans G., Steeghs D., & O'Brien K. 2006b, A&A, 448, 1101
- Karasev, D.I., Lutovinov, A.A., & Burenin, R.A. 2008, Astron. Lett., 34, 753-760; Pisma v Astron.Zh., 34 (2008), 834-841
- Kleinmann, S.G., & Hall, D.N.B. 1986, ApJS, 62, 501
- Maiolino, R., Rieke, G.H., & Rieke, M.J. 1996, AJ, 111, 537
- Malizia, A., Bassani, L., Stephen, J. B., et al. 2005, ApJ, 630, 157
- Martins, F., & Plez, B. 2006, A&A, 457, 637
- Masetti, N., Mason, E., Morelli, L., et al. 2008, A&A, 482, 113
- Mason A.B., Clark J.S., Norton A.J., Negueruela I., Roche P. 2009, arXiv:0907.3876v1
- Messineo M., Figer D., Davies B., Rich M., Valenti E., Kudritzki R.P. 2008, ApJ 683, 155
- Meyer, M.R., Edwards, S., Hinkle, K.H., & Strom S.E. 1998, ApJ, 508, 397
- Negueruela, I., Smith, D.M., Reig, P., et al. 2006a, ESA SP-604, 165
- Negueruela, I., Smith, D.M., Harrison, T.E., & Torrejón, J.M. 2006b, ApJ, 638, 982
- Negueruela, I., Torrejón, J.M., Reig, P., et al. 2008a, AIPC, 1010, 252
- Negueruela, I., Torrejón, J.M., & Reig P. 2009, in Proceedings of the "7th INTEGRAL Workshop - An INTEGRAL View of Compact Objects", Copenhagen, Denmark, 8-11 September 2008, Proceedings of Science.
- Nespoli, E., Fabregat, J., & Mennickent, R.E. 2008, A&A, 486, 911
- Pakull, P., & Parmar, A. 1981, A&A, 102, L1
- Rahoui, F., & Chaty, S. 2008, A&A, 492, 163
- Rahoui, F., Chaty, S., Lagage, P.-O., Pantin, E. 2008, A&A, 484, 801
- Remillard, R., & Smith, D. 2002, ATel 88
- Revnivtsev, M.G., Sunyaev, R.A., Varshalovich, D.A. et al. 2004, Astron. Lett., 30, 382
- Ryter, Ch.E., 1996, Ap&SS, 236, 285
- Schmidt-Kaler, T.H. 1982, in *Landolt-Bornstein New Series, Volume 2b Astronomy and Astrophysics - Stars and Star Clusters*, Eds. K. Schaifers, H.H. Voigt, New York, Springer-Verlag.
- Sguera, V., Barlow, E.J., Bird, A.J., et al. 2005, A&A, 444, 221
- Sguera, V., Bazzano, A., Bird, A.J., et al. 2006, ApJ, 646, 452
- Sguera, V., Hill, A.B., Bird, A.J., et al. 2007, A&A, 467, 249
- Smith, D. M., Heindl, W.A., Markwardt, C.A., et al. 2006, ApJ, 638, 974
- Smith, D.A., Rampy, R.A., Negueruela, I., & Torrejón, J.M. ATel 1268
- Wallace, L.I., Meyer, M.R., Hinkle, K., & Edwards, S. 2000, ApJ, 535, 325
- Walter, R., Zurita-Heras, J., Bassani, L., et al 2006, A&A, 453, 133
- Wegner, W. 1994, MNRAS, 270, 229
- Wegner, W. 2006, MNRAS, 371, 185
- Zurita-Heras, J., & Chaty, S. 2008, A&A, 489, 657
- Zurita-Heras, J., & Chaty, S. 2009, A&A, 493, L1

